


NATIONAL OPEN UNIVERSITY OF NIGERIA
University Village, Plot 91, Cadastral Zone, Nnamdi Azikwe Expressway, Jabi, Abuja

FACULTY OF SCIENCES
DEPARTMENT OF MATHEMATICS

Course Code: MTH381
Course Title: Mathematical Methods III
Credit Unit: 3
Time Allowed: 3 Hours
Total: 70 Marks
Instruction: Answer Question Number One and any Other Four Questions

- 1(a) (i) write the quotient $\frac{1+i}{\sqrt{3}-i}$ in polar form. (2 marks)
(ii) compute $\frac{1+i}{\sqrt{3}-i}$ (1 marks)
(iii) Find all the cube roots of $\sqrt{2} + i\sqrt{2}$ (2 marks)
- 1 (b) (i) Evaluate $|\bar{z}^2 - 2 - (-2 + 2i)|$ (2 marks)
(ii) Evaluate $|\bar{z}^2 - 2| - \sqrt{8}$ (2 marks)
(iii) Let $f(z) = \bar{z}e^{-|z|^2}$. Determine the points at which $f'(z)$ exists, and find $f'(z)$ at these points. (2 marks)
(iv) Find an analytic function f whose imaginary part is given by $e^{-y} \sin x$. (2 marks)
- 1 (c) (i) Use Demoivre's theorem with $n = 4$ to prove that $\cos 4\theta = 8\cos^4\theta - 8\cos^2\theta + 1$
And deduce that $\cos \frac{\pi}{8} = \left(\frac{2+2\sqrt{2}}{4}\right)^{\frac{1}{2}}$. (2 marks)
(ii) Find an analytic function of $z = x + iy$ whose imaginary part is $(y \cos y + x \sin y) e^{px}$ (2 marks)
(iii) Find the radius of convergence of the following Taylor series: (1 marks)
- $$\sum_{n=1}^{\infty} z^n n^{\ln n}$$
- 1 (d) (i) Find the residue at each of the singularities of $f(z) = \frac{e^{z^3}}{z(z+1)}$. The function $f(z)$ has simple poles at $z = 0$ and $z = -1$. Therefore, we have
- $$R[f, 0] = \lim_{z \rightarrow 0} z f(z) = \lim_{z \rightarrow 0} \frac{e^{z^3}}{z(z+1)} = 1$$
- And $R[f, -1] = \lim_{z \rightarrow -1} (z+1)f(z) = \lim_{z \rightarrow -1} \frac{e^{z^3}}{z} = -e^{-1}$ (1 mark)
- (ii) Find the residue of $f(z) = \frac{\cot z}{z^2}$ at $z = 0$. (1 marks)
(iii) Evaluate $I = \int_0^{2\pi} \frac{1}{1+a \sin \theta} d\theta$, $0 < |a| < 1$. (2 marks)



2 (a) (i) Evaluate $\int_0^1 \frac{1}{x^5} dx$ (2 marks)

(ii) show that $\int_{-\infty}^{\infty} \frac{x}{x^3 - a^3} dx = \frac{\pi}{\sqrt{3}a}; a > 0$ (1 mark)

(iii) show that $\int_{-\infty}^{\infty} \frac{\sqrt{x}}{x^3 + 1} dx = \frac{\pi}{3}$ (2 marks)


2 (b) (i) Evaluate $\int_{\gamma} f(z) dz$. If $f(z) = z - 1$ and γ is the curve given by

$$z(t) = t + it^2, \quad 0 \leq t \leq 1. \quad (1 \text{ mark})$$


(ii) Let $f(z) = z - 1$ and $\gamma = \gamma_1 + \gamma_2$ where γ_1 is given by $z_1(t) = t, 0 \leq t \leq 1$ and γ_2 is given by $z_2(t) = 1 + i(t - 1), 1 \leq t \leq 2$, then evaluate $\int_{\gamma} f(z) dz$ (1 mark)


(iii) let γ be given by $z(t) = 2e^{it}, 0 \leq t \leq 2\pi$. Show that $\left| \int_{\gamma} \frac{e^z}{z^2 + 1} dz \right| \leq \frac{4\pi e^2}{3}$ (1 mark)

2 (c) (i) compute the integral $\int_{\gamma} (z^2 - 1) dz$, where γ is the contour in (1 mark)




(ii) compute the integral $\int_{\gamma} \sin z dz$, where γ is the contour (1 mark)






(iii) Compute $\int_{\gamma} \frac{dz}{z}$, where γ is the contour in the given figure (2 marks)





3. (a) Let $\gamma = \gamma_1 + \gamma_2$, where γ_1 and γ_2 respectively are given by $z_1(t) = ti$ and $z_2(t) = t + i$, $t \in [0,1]$. Furthermore, let $f(z) = (y - x) + 3ix^2$. Show that the function f cannot have an antiderivative. (4 marks)

3 (b) (i) Compute $\int_{\gamma} \frac{e^z}{z(z-2)} dz$, where γ is the following contour (4 marks)




(ii) compute $\int_{\gamma} \frac{e^z}{z(z-2)} dz$, where γ is the following contour (4 marks)





4 (a) (i) Compute $\int_{\gamma} \frac{3z+1}{z(z-2)^2} dz$ along the contour γ given in 3b (i) above. **(2 marks)**

(ii) Compute $\int_{\gamma} \frac{\cosh z}{z(z-2)^2} dz$ where the contour γ is given in the figure below **(3 marks)**



(iii) Evaluate the function $f(z) = \int_0^1 e^{-z^2 t} dt$. Let γ be any simple closed contour in the complex plane. Changing the order of integration, we have **(2 marks)**

4 (b) Find the Taylor expansion up to quadratic terms in $x - 2$ and $y - 3$, of $f(x, y) = ye^{xy}$ **(2 marks)**

4 (c) Find and evaluate the maxima and saddle points of the function

$$f(x, y) = xy(x^2 + y^2 - 1) \quad \textbf{(3 marks)}$$

5 (a) (i) by finding $\frac{dl}{dy}$, evaluate the integral **(3 marks)**


$$\int_0^{\infty} \frac{e^{-xy} \sin x}{x} dx$$

(ii) show that the function $1, x, \sin x$ are linearly independent **(2 marks)**

(iii) Find the Laplace transform of the function $f(t) = e^{at}$ **(2 marks)**

5 (b) (i) Evaluate the double integral

$$I = \iint_R x^2 y dx dy \quad \textbf{(2 marks)}$$



where R is the triangular area bounded by the lines $x = 0$, $y = 0$ and $x + y = 1$. Reverse the order of integration and demonstrate that the same results is obtained.

(ii) Find the Laplace transform of $\frac{d^2 f}{dt^2}$ **(3 marks)**

6 (a) In spherical polar coordinates r, θ, ϕ the element of volume for a body that is symmetrical about the polar axis is $dV = 2\pi r^2 \sin \theta dr d\theta$, whilst its element of surface area is $2\pi r \sin \theta [(dr)^2 + r^2 (d\theta)^2]^{\frac{1}{2}}$. A particular surface is defined by $r = 2a \cos \theta$, where a is a constant and $0 \leq \theta \leq \frac{\pi}{2}$. Find its total surface area and the volume it encloses, and hence identify the surface. **(7 marks)**

6 (b) By transforming to cylindrical polar coordinates, evaluate the integral

$$I = \iiint \ln(x^2 + y^2) dx dy dz$$

Over the interior of the conical region $x^2 + y^2 \leq z^2, 0 \leq z \leq 1$ **(5 marks)**